

Review Article

COVID-19: A review of immune-enhancing nutrients and supplements

Hagar F. Forsan ¹, Ayman R. Abdul Rahman², Menna. Safwat ³, Rihan S. Abduljabar⁴, Omnia K. M. Riad³

Abstract

In the era of COVID-19 infection, there is an urgent need to use immune-enhancing nutrients. The frequency of COVID-19 infection is largely affected by an individual's nutritional status. This review summarizes the available nutrients and supplements that help enhance the immune system against infections especially COVID-19. They have antiviral activities, especially against SARS-CoV-2, or even can ameliorate the undesired consequences of COVID-19 by their anti-inflammatory and antioxidant properties. It mentions many nutrients and supplements that aid in fighting or protecting from this pandemic infection such as probiotics, milk including breastfeeding, camel's milk, and supplements including vitamins and minerals. It discusses their role as an antiviral, immunostimulant, or even their role as antioxidant and anti-inflammatory COVID-19 protection. Until our time, the pieces of evidence determining the use of dietary supplements in COVID-19 protection and treatment are insufficient and restricted. So, the use of dietary supplements cannot be used alone to treat COVID-19 but to support the current management of COVID-19.

Authors Affiliation:

¹Dairy Chemistry Research Dept.
Animal Production Research Institute
(APRI), Agriculture Research Center
(ARC), Giza, Egypt.

²Radiology Dept. Armed Forces College
of Medicine, Cairo, Egypt.

³Microbiology and Immunology
Department, Faculty of Pharmacy, Al-
Azhar University, Cairo, Egypt.

⁴Phytochemistry Dept. Scientific
Research Center, Soran University,
Iraq.

Correspondence to: Forsan HF
hagarfathy@pg.cu.edu.eg/
hager.forsan@post.agr.cu.edu.eg/



doi:10.18576/jans/010101

Cite this as:

Forsan HF, Abdul Rahman AR, Safwat
M, Abduljabar RS, Riad OK. COVID-
19: A review of immune-enhancing
nutrients and supplements. *JANS* 2022;
Jan 1(1):1-14.

Received: 27 March 2021

Accepted: 14 June 2021

Copyright © 2022 by the authors.
Licensee Natural Sciences Publishing
Cor, USA.

<http://www.naturalspublishing.com>

Keywords: Coronavirus, Breastfeeding, Immunostimulants, Minerals, Probiotics, Vitamins

Highlights:

- Prompt hydration and a balanced diet are needed to enhance immunity and protect from infectious diseases
- Breastfeeding during COVID-19 infection is recommended after performing the required precautions.
- Probiotics have antiviral and immunomodulatory actions alongside a role in decreasing the cytokine storm with current antiviral therapy.
- Several supplements of vitamins and minerals have an antiviral, antioxidant, anti-inflammatory, and immune-modulatory function.

1. Introduction:

SINCE the present epidemic of the COVID-19 keeps increasing. Now the serious choice is to improve the ability of the immune system to fight against covid-19.

The severity of COVID-19 depends on different factors such as smoking habits, coexisting medical conditions, and the most important factor is the nutritional status which affects immunity. There are a lot of nutrients that help to enhance the immune system ^[1].

2. Nutrition during COVID-19 outbreak:

World health organization (WHO) recommends prompt hydration and eating a balanced diet to fulfill the body's needs of vitamins, minerals, protein, fibers, antioxidants to raise the immune system and shield the body from contagious and chronic illnesses. WHO recommends eating fresh, unprocessed foods and unsaturated fats found in fish, nuts, olive oil, and avocado? WHO warns against eating foods high in salts or sugars and recognizes the need for therapeutic assistance to support the mental health of persons at high risk of COVID-19 infection ^[2].

2.1 Breast milk as the best immunity-enhancing source:

Breast milk is one of the most effective sources for enhancing the immunity of newborns against environmental pathogens. There are major antibodies in breast milk, such as (IgA, IgG, and IgM), transforming growth factor β 2 (TGF- β 2), the soluble cluster of differentiation 14 (sCD14), and antibacterial enzymes such as lysozyme that play significant immune protection role in infant's health ^[3].

There are about 10-fold higher leukocytes in first milk for humans more than mature milk (5 M/ ml). Ten percent of them are NK cells, B cells, and T cells. Neutrophils and macrophages are the main remaining leukocytes, and their numbers decrease with time ^[3]. It also contains massive numbers of innate lymphoid cells that are divided into 1,2,3 lymphoid cells ^[4,5]. These immune cells in a newborn's body destroy pathogens and have a role in shaping an infant's gut microbiota ^[6].

As breastfeeding is a critical and indispensable step in infant life, the WHO and UNICEF are playing an important role in inspiring women to breastfeed during COVID-19 disease - SARS-CoV-2 is unlikely to be passed through breastmilk. Breastfeeding can relieve the possible dangers of COVID-19. Centers for disease control and prevention (CDC) support breastfeeding during COVID-19 infection after performing the required precautions. The precautions to avoid the extent of SARS-CoV-2 infection include handwashing before touching the baby and his tools along with wearing a cloth face covering. If the mother is isolated from her baby, she could express her milk eight to ten times a day using a dedicated pump ^[3].

2.2 Camels' milk as nutritive and therapeutic application

While camels are expected to be a MERS-COV source. The camel's milk can be used as a food of high nutritious and medicinal use. It has excessive

therapeutic uses compared with that of other species such as goat, cow, and buffalo milk. It contains large amounts of essential amino acids (e.g. leucine and phenylalanine), vitamins (e.g., vitamin C and niacin), unsaturated fatty acids, antioxidants, and antimicrobial factors (e.g., lactoperoxidase, lactoferrin, lysozyme, and immunoglobulin G) and secretory immunoglobulin A^[7,8].

Those immunoglobulins cause high antibody titers against rotavirus in an old study^[9]. Other studies suggest that camels' milk has antiviral activity as that of Saltanat and his colleagues who found that camels' milk inhibits the replication of hepatitis B virus and support the response of cellular immunity via correcting the imbalance of Th1/Th2 cytokine and regulating the expression of these cytokines^[10]. Another Egyptian study confirmed that 60 days of drinking camels' milk with peginterferon and ribavirin confers regulation of immunomodulators, inflammatory mediators, antioxidants, antiapoptotic factors for Hepatitis C (HCV) virus infection in patients^[11].

2.3 Probiotic diets and supplements:

Although cytokine referred special effects that are an essential part of the infection response, extreme production of proinflammatory cytokines, or wrong biological context of cytokines, raise the danger of an extensive range of diseases, such as COVID-19.

Inflammation, such as pro-inflammatory and anti-inflammatory cytokines, has a significant role in COVID-19 pathogenesis. Probiotics supplementation can restore innate and adaptive immunity, also modify intestinal microbiota variations which are responsible for immune homeostasis, and the equilibrium balance between Th17 cells and regulatory T-cells (Tregs). It is important to know that probiotics consumption can help to quench the cytokine storm when recombined with suitable antiviral treatments^[12].

Oral probiotics can protect against the systemic and intestinal effects of SARS-CoV-2 infection. Inhaled probiotic provides a direct effect on respiratory immune cells and respiratory epithelium. It can enhance virus clearance and decrease the inflammation caused by this virus. There is an ongoing clinical trial on *Lactobacillus coryneformis* K8 consumption for COVID-19 patients^[13].

2.4 Nutrients and virus-host immunologic responses

From the initial virus-host encounter, innate immune activation, to adaptive immune responses, micronutrients play a role in the host immune response to the virus, as summarized in Fig. 1.

3. Supplementations:

3.1 Omega 3 fatty acids:

Seafood (especially fatty fish) is a rich source of docosahexaenoic acid (DHA), omega-3 (n-3) fatty acids, and eicosatetraenoic acid (EPA). They are also found in many supplements and concentrated pharmaceutical formulations^[15]. Their anti-inflammatory effect was evidenced, especially EPA and docosahexaenoic acid (DHA). The potential effects of n-3 fatty acid supplements are co-therapy for patients afflicted with SARS-CoV-2, including less inflammatory eicosanoids (PG3, TXs, LTs5), non-inflammatory cytokines (TNF-a, IL-1 β),

advanced pro-resolving mediators such as proteins. It has anti-inflammatory, antiplatelet, and potential antiviral effects. The detrimental effects of EPA and DHA are also vulnerable to non-enzymatic oxidation by reactive oxygen species (ROS), possibly toxic oxidation products (MDA, 4-HNE, 4HHE, IsoPs), oxidative stress, and delayed virus clearness^[16].

3.2 Vitamins and Minerals:

Supplementation of minerals and multivitamins including water-soluble and non-water-soluble vitamins that reduce morbidity and mortality rates of severe COVID-19 infection^[17,18].

3.2.1 Vitamins:

3.2.1.1 Vitamin A:

Retinoid supplementation is important for human health. It is largely found in fish oils, animal and fish livers, and fish oil. It is also present in milk and milk products, seafood, shellfish, meats, and poultry, as well as in other food sources such as dark leafy plants, algae, red, yellow vegetables, and tubers^[19].

Vitamin A ensures epithelial tissue barrier stability and normal separation. Retinoid has a role in immune function by regulating NK cells and encouraging the phagocytic oxidative burst activities of macrophages^[20,21], development, and differentiation of the Th1/Th2^[22]. A recent study indicates that vitamin A is used in the treatment of COVID-19 depending on bioinformatics analysis, they confirmed that vitamin A enriches the immune reactions and has an anti-inflammatory function and decreases reductive oxygen species. They also identified IL10, MAPK1, EGFR, MAPK14, ICAM1, CAT, and PRKCB as the main targets against COVID-19^[23].

3.2.1.2 Vitamin B6:

Nature is full of hundreds of carotenoids but there are few amounts found in human tissues. Beta-carotene is a pro-vitamin A in the diet. Almost 50 carotenoids have pro-vitamin A activity, but beta-carotene is the most powerful and widely distributed carotenoid in plant species.

Diet and nutrition such as milk and soya beans have a positive impact on immunity, especially against COVID-19. It is documented that pyridoxine (the active form of vitamin B6) is depleted gradually during inflammation. As COVID-19 causes many inflammatory reactions, it may be recommended to administer vitamin B6^[24].

3.2.1.3 Vitamin C:

Vitamin C can't be produced in the human body. It motivates absorption of non-heme iron from the intestine and moderates transport and storage of iron that helps in anemia prevention^[25]. It has an essential role in epithelial barrier integrity^[26] and immune cell activities including the production of antibodies, T-cells proliferation, and differentiation^[27].

It supports the production of antimicrobial agents such as IFN γ and increases serum complement proteins^[28]. It's an antioxidant that keeps the

intracellular reductive-oxidative homeostasis ^[29]. The increased oxidative stress in COVID-19 patients caused by free radicals and cytokine's rapid release during a cytokine storm makes an urgent need to add antioxidants in the treatment protocols for COVID-19, vitamin C is the most suggested antioxidant ^[30].

Some clinical trials were done to determine the impact of using high doses of vitamin C (oral or intravenous administration). Staying in the intensive care unit (ICU) decreased by 7.8% after 200 mg/kg body weight/day consumption [31]. Some studies have shown that oral vitamin C decreases the risk of infection with viruses ^[30]. Analysis of 17 COVID-19 patients found that the use of high-dose I.V. Vitamin C can be used in patients with moderate-severe illnesses ^[31].

In the last trial to assess the use of high-dose of vitamin C (intravenous) for COVID-19; they found that vitamin C may be useful for oxygenation of severe COVID-19. They revealed that intravenous vitamin C in high doses does not affect patients who need invasive mechanical ventilation ^[32]. Thomas and her colleagues reported that the use of high-dose of vitamin C, zinc gluconate, or a combination of them have an insignificant decrease in the duration of symptoms ^[33]. Assessing the COVID-19 patients for their vitamins and minerals deficiencies is an indispensable step in the correct management of COVID-19 ^[34].




Fig.1. The immunomodulatory activity of different dietary substances in different stages of the host immune response to the virus ^[14]

3.2.1.4 Vitamin D:

Vitamin D supplementation is an important source for the prevention of cancer, osteoarthritis progression, multiple sclerosis, and hypertension ^[35]. Milk and dairy products are the best vitamin D sources. The general actions and metabolism of vitamin D are well recognized. Vitamin D₃ is formed by the effects of UVB radiation on 7-dehydrocholesterol, followed by a thermal reaction in the skin. It helps to increase immune barrier integrity ^[35], antimicrobial peptides production ^[36] and it supports macrophages, monocytes, and dendritic cell functions ^[37].

It regulates oxidative burst potential^[38] and IFN γ inhibition^[39]. Vitamin D is confirmed to improve innate immunity by activating toll-like receptors (TLRs) or the number of β -defensins and cathelicidins.

Vitamin D's immunostimulant effect is not limited to intrinsic immunity, it affects adaptive immunity, and it modulates the function of T cells. Supplementation with vitamin D could help in combating the SARS-CoV-2 infection^[40].

3.2.1.5 Vitamin E:

It is a vital antioxidant, that prevents oxidative damage to the cell membranes and rises respiratory epithelial barriers stability^[41]. It may increase the cytotoxic activity of NK cells and macrophages, decrease the production of prostaglandin E2, and change IFN- γ and interleukin 2 development^[42].

It helps T-cells lymphocytes proliferation and mediates optimize Th1 response, supports Th2 response suppression, encourages effective immune synapses of vitamin E protection including Th cells and helps to increase the amount of antigen-experienced T-cell memory^[43]. Almonds and peanuts are vitamin E sources.

3.2.2 Minerals:

3.2.2.1 Zinc:

Zinc (Zn) is a significant micronutrient for human beings. Cereals including wheat are good sources of it^[44]. Zinc deficiency is a predisposing factor for the progression of COVID-19 infection, as its deficiency is correlated with reduced breathing epithelial integrity, increased viral entry, and imbalanced immune and redox system^[12]. It modifies the biological and physiological roles of approximately 2,000 enzymes and 750 transcription factors involved in the development of immunity^[45].

Zn can increase the anticancer bioactivity, promote the production and differentiation of essential immune cells^[46]. It is correlated with the activities of complement proteins, the IFN γ creation, T- cells, and antibody production essentially the immunoglobulin G^[29]. It also has a direct antiviral property that prevents the RNA-dependent RNA polymerase enzyme^[29]. So, it is strongly recommended to add zinc in the treatment protocols of COVID-19 either alone or combined with other strategies^[47]. There is a recent study that mentioned that adding zinc to the treatment protocol of COVID- 19 containing low doses of hydroxychloroquine is associated with significantly lower hospitalization^[48].

3.2.2.2. Selenium:

Selenium is an important micronutrient in the living systems^[49]. It plays a necessary role in enhancing antioxidant defense capacity, improve immunity, metabolism of thyroid hormone, and function reproduction^[50]. So, it is relevant to develop new food sources of selenium^[50]. A recent study reported that COVID-19 mortality is increased with selenium deficiency^[51]. It helps in the formation of killer cells, T-cells, and the production of antibodies^[52]. Cereals and dairy products are selenium sources.

3.2.2.3 Magnesium:

Cereals and vegetables are healthy sources of magnesium, while milk and dairy products are considered better sources^[53]. It is incorporated in the synthesis of nucleic acid, transcription of DNA, encouragement of leukocytes, macrophage antigen binding, and apoptotic directive^[39]. It influences cell-mediated immunity as well as humoral adaptive immunity^[21]. It keeps DNA from oxidative damage and inhibits the production of superoxide anions at high concentrations^[52].

3.2.2.4 Copper:

For humans, copper (Cu) is an important ingredient. It is present in different grains as wheat^[54]. During viral infection, copper (Cu) is a vital micronutrient. Cu improves the function of extreme immune cells, including T helper cells, B cells, NK neutrophils, and macrophages. These blood cells are responsible for infectious microbes being killed. Cu-deficient humans play a significant benefit in infections because of the reduced number and function of these blood cells. In addition, Cu is capable of successfully destroying various viruses, such as SARS-CoV-2.^[55] Table 1 shows the mechanism of fighting COVID-19 by different dietary supplements.

Table 1: Beneficial dietary supplements in COVID-19 protection and treatment:

| Dietary supplements | Role of dietary supplement in fighting COVID-19 |
|---------------------|--|
| Omega 3 FAs | <ul style="list-style-type: none"> It has anti-inflammatory and antiviral activity^[16] |
| Vitamin A | <ul style="list-style-type: none"> It ensures epithelial tissue barrier stability^[20, 21]. It stimulates innate and adaptive immunity^[21, 22]. |
| Vitamin B6 | <ul style="list-style-type: none"> It has anti-inflammatory activity^[23] Vitamin B6 is recommended during COVID-19 to compensate for the pyridoxine depletion which happens during inflammation^[24]. |
| Vitamin C | <ul style="list-style-type: none"> It increases epithelial barrier integrity^[27]. Stimulate innate and adaptive immune responses^[27]. It has potent antioxidant activity^[29]. It supports the production of IFN γ and serum complement proteins^[28]. |
| Vitamin D | <ul style="list-style-type: none"> It increases immune barrier integrity^[35]. It increases antimicrobial peptide production^[35]. It supports innate and adaptive immune responses^[40]. |
| Vitamin E | <ul style="list-style-type: none"> It has antioxidant activity^[41]. It supports adaptive immune responses^[43]. |
| Zinc | <ul style="list-style-type: none"> Zn promotes the production and differentiation of essential immune cells^[46] and increases the activity of complement proteins, T- cells, and antibody production^[29]. |
| Selenium | <ul style="list-style-type: none"> It has a direct antiviral property^[29]. It has antioxidant activity^[50]. It improves immunity, the formation of killer cells, T-cells, and the production of antibodies^{[50], [52]}. |
| Magnesium | <ul style="list-style-type: none"> It supports cell-mediated and humoral adaptive immunity [21]. It has antioxidant activity^[52]. |
| Copper | <ul style="list-style-type: none"> Cu improves the function of extreme immune cells, including T helper cells, B cells, NK neutrophils, and macrophages. Cu can destroy various viruses, such as SARS-CoV-2^[55]. |

4. Additional potential immunomodulatory agents for COVID-19 management:

4.1 N-acetylcysteine (NAC):

NAC can treat Glucose 6-phosphate dehydrogenase (G6PD) deficiency and prevent glutathione depletion caused by COVID-19 infection. A recent study is conducted on a patient with G6PD deficiency with COVID-19 infection, they demonstrated that intravenous (IV) NAC prevents hemolysis, the elevation of C-reactive protein (CRP), liver enzymes, and ferritin. Intravenous NAC facilitates full retrieval of G6PD deficient COVID-19 patients. NAC may be involved in viral infection blockage and cytokine storm alleviation^[56]. NAC is a pioneer of decreased glutathione. Due to its acceptability, this pleiotropic drug has been suggested not only as a mucolytic agent but also as a defensive/ therapeutic agent in a diversity of patients relating to reduction and oxidative stress. The oral administration of N- Acetyl-L- cysteine is probable reducing the risk of COVID-19 emerges. Furthermore, high-dose intravenous N- Acetyl-L-cysteine can play an important role in COVID-19 cases treatment^[57].

5. The use of dietary supplements in light of the levels of evidence-based medicine:

It is worth saying that the evidence which determines the use of food supplements in COVID-19 protection and treatment is insufficient and restricted^[58] to do well-designed systematic reviews and meta-analysis from randomized controlled trials and reach accurate advice. For example, vitamin D, one of the most studied vitamins regarding its deficiency with the severity of COVID-19, there are only 13 registered clinical trials, only 4 of them are completed^[59]. Most studies discuss the use of supplements through their previous data regarding protection from respiratory viral infections and opposing the mechanism of SARS-CoV-2 pathogenesis. By the time, studies that are especially targeting COVID-19 are increasing facilitating the accessibility to accurate information.

In a large study which is based on an application survey among UK people and included 372720 persons to know the correlation between dietary supplements administration and being positive to SARS-CoV-2, they found that vitamin D, probiotics, omega-3 fatty acids, and multivitamins which include vitamin A, D, E, zinc, and selenium decrease the risk of SARS-CoV-2 infection in females only. They found that vitamin C, zinc, and garlic have no protective effect from SARS-CoV-2 infection. A large number of randomized controlled trials of dietary supplements are needed^[60].

6. Conclusion:

There is an urgent need to find different strategies to control the highly virulent SARS-CoV-2 virus which affects most humans' organs^[61]. Nutrients can show an effective role in the prevention of SARS-CoV-2 infection. Camel's milk, breastfeeding, probiotics, and supplements (i.e. vitamin D, vitamin C, vitamin A, zinc, selenium, magnesium, and copper) have an obvious role in immune modulation and help to fight this pandemic infection. Some supplements may have antiviral properties such as zinc, vitamin C, vitamin D, and probiotics.

Vitamin C and zinc increase respiratory epithelial integrity. Vitamin C, selenium, and magnesium have strong antioxidant activities which protect from the effect of SARS-CoV-2 on the human being. Fatal cytokine storm can be reduced by the anti-inflammatory actions of omega-3, probiotics, vitamin A and vitamin B6 with the available antivirals. The general recommendation is to consume dietary supplements (i.e. vitamins C and D, zinc, and selenium) that should be administered to individuals with or at risk of respiratory viral infections or in whom deficiency is detected.

List of abbreviations:

CDC: Centers for Disease Control

COVID-19: Coronavirus Disease 2019

Cu: Copper

DHA: Docosahexaenoic Acid

EPA: Eicosapentaenoic Acid

Fe: Iron

G6PD: Glucose 6-Phosphate Dehydrogenase

HCV: Hepatitis C Virus

ICU: The Intensive Care Unit IV: Intravenous

M: Million

NAC: N-acetylcysteine

NK: Natural Killer

n-3: omega-3

ROS: Reactive Oxygen Species

Scd14: Soluble Cluster of Differentiation 14

TGF- β 2: Transforming Growth Factor β 2 TLRs: Toll-like Receptors

WHO: World health organization

Zn: Zinc

Conflict of interest

The authors declare that they have no conflict of interest regarding this study.

Author contribution

All authors have contributed in writing the manuscript. HFF & OKR participated in conceptualization. OKR was responsible for revising and editing the manuscript.

Funding: There is no funding source for this study.

References:

- ⁽¹⁾ Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of Coronavirus Disease in China. *N Engl J Med.* 2020 Apr 30;382(18):1708-20. doi: 10.1056/NEJMoa2002032 [Google Scholar] [PubMed].
- ⁽²⁾ WHO. WHO Director-General's opening remarks at the media briefing on COVID-19 2020 [Available from: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19--11-march-2020>] (accessed Feb 29, 2020).
- ⁽³⁾ CDC. If you are pregnant, breastfeeding, or caring for young children 2020 [Available from: <https://www.uncmedicalcenter.org/app/files/public/aed467f6-1ccf-46c1-992f-43c3cd714e47/pdf-med-ctr-whic-if-you-are-pregnant-breastfeeding-or-caring-for-young-children.pdf>] (accessed March 13, 2020).
- ⁽⁴⁾ Baban B, Malik A, Bhatia J, Yu JC. Presence and profile of innate lymphoid cells in human breast milk. *JAMA Pediatr.* 2018;172(6):594-6. doi: 10.1001/jamapediatrics.2018.0148. [Google Scholar] [PubMed].
- ⁽⁵⁾ Palmeira P, Carneiro-Sampaio M. Immunology of breast milk. *Rev Assoc Med Bras* (1992). 2016;62(6):584-93. doi: 10.1590/1806-9282.62.06.584 [Google Scholar] [PubMed].
- ⁽⁶⁾ Eberl G, Colonna M, Di Santo JP, McKenzie AN. Innate lymphoid cells. Innate lymphoid cells: a new paradigm in immunology. *Science.* 2015;348(6237):aaa6566. doi: 10.1126/science.aaa6566 [Google Scholar] [PubMed].
- ⁽⁷⁾ Reusken CB, Farag EA, Jonges M, Godeke GJ, El-Sayed AM, Pas SD, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) RNA and neutralising antibodies in milk collected according to local customs from dromedary camels, Qatar, April 2014. *Euro Surveill.* 2014;19(23) doi: 10.2807/1560-7917.es2014.19.23.20829. [Google Scholar] [PubMed].
- ⁽⁸⁾ Malik YS, Sircar S, Bhat S, Sharun K, Dhama K, Dadar M, et al. Emerging novel coronavirus (2019-nCoV)-current scenario, evolutionary perspective based on genome analysis and recent developments. *Veterinary Quarterly.* 2020;40(1):68-76 doi: 10.1080/01652176.2020.1727993 [Google Scholar] [PubMed].
- ⁽⁹⁾ El Agamy EI, Ruppanner R, Ismail A, Champagne CP, Assaf R. Antibacterial and antiviral activity of camel milk protective proteins. *J Dairy Res.* 1992;59(2):169-75 doi: 10.1017/s0022029900030417 [Google Scholar] [PubMed].
- ⁽¹⁰⁾ Saltanat H, Li H, Xu Y, Wang J, Liu F, Geng XH. The influences of camel milk on the immune response of chronic hepatitis B patients. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi.* 2009;25(5):431-3. Chinese (abstract) [Google Scholar] [PubMed].
- ⁽¹¹⁾ Mohamed WA, Schaalán MF, El-Abhar HS. Camel Milk: Potential Utility as an Adjunctive Therapy to Peg-IFN/RBV in HCV-4 Infected Patients in Egypt. *Nutr Cancer.* 2015;67(8):1305-13 doi: 10.1080/01635581.2015.1087041 [Google Scholar] [PubMed].
- ⁽¹²⁾ Romano L, Bilotta F, Dauri M, Macheda S, Pujia A, De Santis GL, et al. Short Report-Medical nutrition therapy for critically ill patients with COVID-19. *Eur Rev Med Pharmacol Sci.* 2020;24(7):4035-9. doi: 10.26355/eurrev_202004_20874 [Google Scholar] [PubMed].
- ⁽¹³⁾ Infusino F, Marazzato M, Mancone M, Fedele F, Mastroianni CM, Severino P, et al. Diet Supplementation, Probiotics, and Nutraceuticals in SARS-CoV-2 Infection: A Scoping

- Review. *Nutrients*. 2020 Jun 8;12(6):1718. doi: 10.3390/nu12061718 [[Google Scholar](#)] [[PubMed](#)].
- ⁽¹⁴⁾ Gasmi A, Tippairote T, Mujawdiya PK, Peana M, Menzel A, Dadar M, et al. Micronutrients as immunomodulatory tools for COVID-19 management. *Clin Immunol*. 2020 Nov;220:108545. doi: 10.1016/j.clim.2020.108545 [[Google Scholar](#)] [[PubMed](#)].
- ⁽¹⁵⁾ Innes JK, Calder PC. Marine Omega-3 (N-3) Fatty Acids for Cardiovascular Health: An Update for 2020. *Int J Mol Sci*. 2020; 18;21(4):1362. doi: 10.3390/ijms21041362 [[Google Scholar](#)] [[PubMed](#)].
- ⁽¹⁶⁾ Rogero MM, Leao MC, Santana TM, Pimentel M, Carlini GCG, da Silveira TFF, et al. Potential benefits and risks of omega-3 fatty acids supplementation to patients with COVID-19. *Free Radic Biol Med*. 2020;156:190-199. doi: 10.1016/j.freeradbiomed.2020.07.005 [[Google Scholar](#)] [[PubMed](#)].
- ⁽¹⁷⁾ Beigmohammadi MT, Bitarafan S, Hoseindokht A, Abdollahi A, Amoozadeh L, Mahmoodi Ali Abadi M, et al. Impact of vitamins A, B, C, D, and E supplementation on improvement and mortality rate in ICU patients with coronavirus-19: a structured summary of a study protocol for a randomized controlled trial. *Trials*. 2020;21(1):614. doi: 10.1186/s13063-020-04547-0 [[Google Scholar](#)] [[PubMed](#)].
- ⁽¹⁸⁾ Wishart K. Increased Micronutrient Requirements during Physiologically Demanding Situations: Review of the Current Evidence. *Vitam Miner* 2017;6(3):1-16. doi:10.4172/2376-1318.1000166 [[Google Scholar](#)]
- ⁽¹⁹⁾ Booth SL, Johns T, Kuhnlein HV. Natural Food Sources of Vitamin A and Provitamin. *Food Nutr Bull* 1992;14(1):1-15. doi.org/10.1177/156482659201400115. [[Google Scholar](#)].
- ⁽²⁰⁾ Huang Z, Liu Y, Qi G, Brand D, Zheng SG. Role of Vitamin A in the Immune System. *J Clin Med*. 2018;7(9):258. doi: 10.3390/jcm7090258. [[Google Scholar](#)] [[PubMed](#)].
- ⁽²¹⁾ Chaigne-Delalande B, Li FY, O'Connor GM, Lukacs MJ, Jiang P, Zheng L, et al. Mg²⁺ regulates cytotoxic functions of NK and CD8 T cells in chronic EBV infection through NKG2D. *Science*. 2013;12;341(6142):186-91. doi: 10.1126/science.1240094 [[Google Scholar](#)] [[PubMed](#)].
- ⁽²²⁾ Maggini S, Pierre A, Calder PC. Immune function and micronutrient requirements change over the life course. *Nutrients*. 2018;10(10):1531. doi: 10.3390/nu10101531. [[Google Scholar](#)] [[PubMed](#)].
- ⁽²³⁾ Li R, Wu K, Li Y, Liang X, Tse WKF, Yang L, et al. Revealing the targets and mechanisms of vitamin A in the treatment of COVID-19. *Ageing (Albany NY)*. 2020;12(15):15784-96. doi: 10.18632/aging.103888 [[Google Scholar](#)] [[PubMed](#)].
- ⁽²⁴⁾ Paul L, Ueland PM, Selhub J. Mechanistic perspective on the relationship between pyridoxal 5'-phosphate and inflammation. *Nutr Rev* 2013;71(4):239-44. doi: 10.1111/nure.12014. [[Google Scholar](#)] [[PubMed](#)].
- ⁽²⁵⁾ Hemilä H, Chalker E. Commentary: Vitamin C supplementation for prevention and treatment of pneumonia. *Front Med (Lausanne)*. 2021;20;7:595988. doi: 10.3389/fmed.2020.595988. [[Google Scholar](#)] [[PubMed](#)].
- ⁽²⁶⁾ Carr AC, Maggini S. Vitamin C and Immune Function. *Nutrients*. 2017; 3;9(11):1211. doi: 10.3390/nu9111211. [[Google Scholar](#)] [[PubMed](#)].

- ⁽²⁷⁾ Hiedra R, Lo KB, Elbashesheh M, Gul F, Wright RM, Albano J, et al. The use of IV vitamin C for patients with COVID-19: a case series. *Expert Rev Anti Infect Ther.* 2020;18(12):1259-61. doi: 10.1080/14787210.2020.1794819. [Google Scholar] [PubMed].
- ⁽²⁸⁾ Bozonet SM, Carr AC. The Role of Physiological Vitamin C Concentrations on Key Functions of Neutrophils Isolated from Healthy Individuals. *Nutrients.* 2019 Jun 17;11(6):1363. doi: 10.3390/nu11061363. [Google Scholar] [PubMed].
- ⁽²⁹⁾ Wintergerst ES, Maggini S, Hornig DH. Immune-enhancing role of vitamin C and zinc and effect on clinical conditions. *Ann Nutr Metab.* 2006;50(2):85-94. doi: 10.1159/000090495 [Google Scholar] [PubMed].
- ⁽³⁰⁾ Cheng RZ. Can early and high intravenous dose of vitamin C prevent and treat coronavirus disease 2019 (COVID-19)? *Med Drug Discov.* 2020 Mar;5:100028. doi: 10.1016/j.medidd.2020.100028 [Google Scholar] [PubMed].
- ⁽³¹⁾ Hemilä H, Chalker E. Vitamin C Can Shorten the Length of Stay in the ICU: A Meta-Analysis. *Nutrients* 2019;11(4):708. doi: 10.3390/nu11040708 [Google Scholar] [PubMed].
- ⁽³²⁾ Zhang J, Rao X, Li Y, Zhu Y, Liu F, Guo G, et al. Pilot trial of high-dose vitamin C in critically ill COVID-19 patients. *Ann Intensive Care.* 2021;9;11(1):5. doi: 10.1186/s13613-020-00792-3 [Google Scholar] [PubMed].
- ⁽³³⁾ Thomas S, Patel D, Bittel B, Wolski K, Wang Q, Kumar A, et al. Effect of High-Dose Zinc and Ascorbic Acid Supplementation vs Usual Care on Symptom Length and Reduction Among Ambulatory Patients With SARS-CoV-2 Infection: The COVID A to Z Randomized Clinical Trial. *JAMA Netw Open.* 2021 Feb 1;4(2):e210369. doi: 10.1001/jamanetworkopen.2021.0369 [Google Scholar] [PubMed].
- ⁽³⁴⁾ Carr AC. Micronutrient status of COVID-19 patients: a critical consideration. *Critical Care.* 2020;16;24(1):1-2. doi: 10.1186/s13054-020-03085-0. [Google Scholar] [PubMed].
- ⁽³⁵⁾ Vieth R. Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr.* 1999;69(5):842-56. doi: 10.1093/ajcn/69.5.842 [Google Scholar] [PubMed].
- ⁽³⁶⁾ WHO/FAO Joint. *Vitamin and mineral requirements in human nutrition.* 2nd ed. Printed in China by Sun Fung 2004. [Available from: <https://apps.who.int/iris/handle/10665/42716>].
- ⁽³⁷⁾ Biesalski HK. Nutrition meets the microbiome: micronutrients and the microbiota. *Ann N Y Acad Sci.* 2016;1372(1):53-64. doi: 10.1111/nyas.13145. [Google Scholar] [PubMed].
- ⁽³⁸⁾ Gombart AF. The vitamin D-antimicrobial peptide pathway and its role in protection against infection. *Future Microbiol.* 2009;4(9):1151-65. doi: 10.2217/fmb.09.87. [Google Scholar] [PubMed].
- ⁽³⁹⁾ Petrovic J, Stanic D, Dmitrasinovic G, Plecas-Solarovic B, Ignjatovic S, Batinic B, et al. Magnesium Supplementation Diminishes Peripheral Blood Lymphocyte DNA Oxidative Damage in Athletes and Sedentary Young Man. *Oxid Med Cell Longev.* 2016;2016:2019643. doi: 10.1155/2016/2019643. [Google Scholar] [PubMed].
- ⁽⁴⁰⁾ Lin Z, Li W. The Roles of Vitamin D and Its Analogs in Inflammatory Diseases. *Curr Top Med Chem.* 2016;16(11):1242-61. doi: 10.2174/1568026615666150915111557. [Google Scholar] [PubMed].

- ⁽⁴¹⁾ Lee G, Han S. The Role of Vitamin E in Immunity. *Nutrients*. 2018;1;10(11):1614. doi: 10.3390/nu10111614. [Google Scholar] [PubMed].
- ⁽⁴²⁾ Wu D, Lewis ED, Pae M, Meydani SN. Nutritional Modulation of Immune Function: Analysis of Evidence, Mechanisms, and Clinical Relevance. *Front Immunol*. 2019;15;9:3160. doi: 10.3389/fimmu.2018.03160. [Google Scholar] [PubMed].
- ⁽⁴³⁾ Han SN, Adolfsson O, Lee CK, Prolla TA, Ordovas J, Meydani SN. Vitamin E and gene expression in immune cells. *Ann N Y Acad Sci*. 2004;1031:96-101. doi: 10.1196/annals.1331.010. [Google Scholar] [PubMed].
- ⁽⁴⁴⁾ Khokhar JS, King J, King IP, Young SD, Foulkes MJ, Silva JD, et al. Novel sources of variation in grain Zinc (Zn) concentration in bread wheat germplasm derived from Watkins landraces. *PLoS one*. 2020;28;15(2):0229107. doi: 10.1371/journal.pone.0229107. [Google Scholar] [PubMed].
- ⁽⁴⁵⁾ Read SA, Obeid S, Ahlenstiel C, Ahlenstiel G. The role of zinc in antiviral immunity. *Adv Nutr*. 2019;1;10(4):696-710. doi: 10.1093/advances/nmz013. [Google Scholar] [PubMed].
- ⁽⁴⁶⁾ Ghaffari H, Tavakoli A, Moradi A, Tabarraei A, Bokharaei-Salim F, Zahmatkeshan M, et al. Inhibition of H1N1 influenza virus infection by zinc oxide nanoparticles: another emerging application of nanomedicine. *J Biomed Sci*. 2019;10;26(1):1-10. doi: 10.1186/s12929-019-0563-4. [Google Scholar] [PubMed].
- ⁽⁴⁷⁾ Cluver L, Jamie ML, Lorraine Sherr, Inge Wessels, Etienne Krug, Sabine Rakotomalala, et al. Parenting in a time of COVID-19. *Lancet*. 2020;11;395(10231):e64. doi: 10.1016/S0140-6736(20)30736-4. [Google Scholar] [PubMed].
- ⁽⁴⁸⁾ Derwand R, Scholz M. Does zinc supplementation enhance the clinical efficacy of chloroquine/ hydroxychloroquine to win today's battle against COVID-19? *Med Hypotheses*. 2020 Sep;142:109815. doi: 10.1016/j.mehy.2020.109815. [Google Scholar] [PubMed].
- ⁽⁴⁹⁾ Glotova IA, Galochkina NA. Development, properties and sanitary-hygienic assessment of selenium-containing cereal food additives. *IOP Conf. Ser.: Earth Environ. Sci*. 2020;422; 01208. doi.org/10.1088/1755-1315/422/1/012085. [Google Scholar]
- ⁽⁵⁰⁾ Zhang S, Xie Y, Li M, Yang H, Li S, Li J, et al. Effects of Different Selenium Sources on Meat Quality and Shelf Life of Fattening Pigs. *Animals (Basel)*. 2020;3;10(4):615. doi: 10.3390/ani10040615. [Google Scholar] [PubMed].
- ⁽⁵¹⁾ Moghaddam A, Heller RA, Sun Q, Seelig J, Cherkezov A, Seibert L, et al. Selenium Deficiency Is Associated with Mortality Risk from COVID-19. *Nutrients*. 2020; 16;12(7):2098. doi: 10.3390/nu12072098. [Google Scholar] [PubMed].
- ⁽⁵²⁾ The Institute for Functional Medicine. IFM, Calendar P. Boosting Immunity: Functional Medicine Tips on Prevention & Immunity Boosting During the COVID-19 (Coronavirus) Outbreak. 2020; [Available from: <https://www.ifm.org/news-insights/boosting-immunity-functional-medicine-tips-prevention-immunity-boosting-covid-19-coronavirus-outbreak/>].
- ⁽⁵³⁾ Cuadrado-Soto E, López-Sobaler AM, Jiménez-Ortega AI, Aparicio A, Bermejo LM, Hernández-Ruiz Á, et al. Usual Dietary Intake, Nutritional Adequacy and Food Sources of Calcium, Phosphorus, Magnesium and Vitamin D of Spanish Children Aged One to <10 Years. Findings from the EsNuPI Study. *Nutrients*. 2020;16;12(6):1787. doi: 10.3390/nu12061787. [Google Scholar] [PubMed].

- ⁶⁴ Sharma S, Nagpal AK, Kaur I. Heavy metal contamination in soil, food crops and associated health risks for residents of Ropar wetland, Punjab, India and its environs. *Food Chem.* 2018;30; 255:15-22. doi: 10.1016/j.foodchem.2018.02.037. [[Google Scholar](#)] [[PubMed](#)].
- ⁶⁵ Raha S, Mallick R, Basak S, Duttaroy A. Is copper beneficial for COVID-19 patients? *Med Hypotheses.* 2020;142:109814. doi: 10.1016/j.mehy.2020.109814. [[Google Scholar](#)] [[PubMed](#)].
- ⁶⁶ Ibrahim H, Perl A, Smith D, Lewis T, Kon Z, Goldenberg R, et al. Therapeutic blockade of inflammation in severe COVID-19 infection with intravenous N- acetylcysteine. *Clin Immunol* 2020;1:219-108544. doi: 10.1016/j.clim.2020.108544. [[Google Scholar](#)] [[PubMed](#)].
- ⁶⁷ De Flora S, Balansky R, La Maestra S. Rationale for the use of N-acetylcysteine in both prevention and adjuvant therapy of COVID-19. *FASEB J.* 2020;34(10):13185-93. doi: 10.1096/fj.202001807 [[Google Scholar](#)] [[PubMed](#)].
- ⁶⁸ Adams KK, Baker WL, Sobieraj DM. Myth Busters: Dietary Supplements and COVID-19. *Ann Pharmacother.* 2020 Aug; 54 (8): 820-826. doi: 10.1177/1060028020928052. [[Google Scholar](#)] [[PubMed](#)].
- ⁶⁹ U. S. National library of medicine. users of the COVID-19 Symptom Study app. *BMJ Nutrition, Prevention & Health* 2021;bmjnph-2021-000250. [Available from: https://clinicaltrials.gov/ct2/results?cond=COVID-19+AND+%22Vitamin+D+Deficiency%22&Search=Apply&recrs=e&age_v=&gndr=&type=&rslt=].
- ⁶⁰ Louca P, Murray B, Klaser K, Graham MS, Mazidi M, Leeming ER. et al. Modest effects of dietary supplements during the COVID-19 pandemic: insights from 445 850 users of the COVID-19 Symptom Study app. *BMJ Nutr Prev Health.* 2021 Apr 19;4(1):149-157. doi: 10.1136/bmjnph-2021-000250. [[Google Scholar](#)] [[PubMed](#)].
- ⁶¹ Safwat M, Forsan H, Abdul-Rahman A, Abduljabar R, Riad O. COVID-19; immunology, pathology, severity and immunosuppressants. *Azhar Int J Pharm Med Sci.* 2021;1(1): 1-12. doi: 10.21608/aijpm.2021.50103.1005. [[Google Scholar](#)].