

Dynamics of the Rational Difference Equation

M. A. El-Moneam^{1,*} and E. M. E. Zayed²

¹ Mathematics Department, Faculty of Science and Arts in Farasan, Jazan University, Kingdom of Saudi Arabia

² Mathematics Department, Faculty of Science, Zagazig University, Zagazig, Egypt

Received: 8 Dec. 2013, Revised: 17 Mar. 2014, Accepted: 18 Mar. 2014

Published online: 1 May 2014

Abstract: In this article, we study the periodicity, the boundedness and the global stability of the positive solutions of the following nonlinear difference equation

$$x_{n+1} = Ax_n + Bx_{n-k} + Cx_{n-l} + Dx_{n-\sigma} + \frac{bx_{n-k} + hx_{n-l}}{dx_{n-k} + ex_{n-l}}, \quad n = 0, 1, 2, \dots,$$

where the coefficients $A, B, C, D, b, d, e, h \in (0, \infty)$, while k, l and σ are positive integers. The initial conditions $x_{-\sigma}, \dots, x_{-l}, \dots, x_{-k}, \dots, x_{-1}, x_0$ are arbitrary positive real numbers such that $k < l < \sigma$. We will prove that the equilibrium points of this equation are locally asymptotically stable, global attractor and hence they are global stability. This equation will have (or have not) prime period two solution under suitable conditions on these coefficients. The solutions of this equation will be proved to be bounded. Some numerical examples will be given to illustrate our results.

Keywords: Difference equations, prime period two solution, boundedness character, locally asymptotically stable, global attractor, global stability.

1 Introduction

The topic of difference equations is interesting and attractive to many mathematicians working in this field. It is a fertile research area. Many real life phenomena are modelling using these equations. Examples from economy, biology, etc. may be obtained in [11, 16, 17]. The study of some properties of these equations via the global attractivity, the boundedness and the periodicity of these equations is of great interest. For example, in the articles [2–10] [12–17] closely related global convergence results were obtained which can be applied to nonlinear difference equations in proving that every solution of these equations converges to a period two solution. For other closely related results, (see [18–25], [27–29]) and the references cited therein.

The objective of this article is to investigate some qualitative behavior of the solutions of the nonlinear difference equation

$$x_{n+1} = Ax_n + Bx_{n-k} + Cx_{n-l} + Dx_{n-\sigma} + \frac{bx_{n-k} + hx_{n-l}}{dx_{n-k} + ex_{n-l}}, \quad n = 0, 1, 2, \dots \tag{1}$$

where the coefficients $A, B, C, D, b, d, e, h \in (0, \infty)$, while k, l and σ are positive integers. The initial conditions $x_{-\sigma}, \dots, x_{-l}, \dots, x_{-k}, \dots, x_{-1}, x_0$ are arbitrary positive real numbers such that $k < l < \sigma$. Note that the special cases of Eq.(1) have been discussed in [1, 26, 30, 31].

Our interest now is to study the behavior of the solutions of Eq.(1) in its general form. For the related work, (see [32–37]). Let us now recall some well known results [11] which will be useful in the sequel.

Definition 1. Consider a difference equation in the form

$$x_{n+1} = F(x_n, x_{n-k}, x_{n-l}, x_{n-\sigma}), \quad n = 0, 1, 2, \dots \tag{2}$$

where F is a continuous function, while k and l are positive integers such that $k < l < \sigma$. An equilibrium point \tilde{x} of this equation is a point that satisfies the condition $\tilde{x} = F(\tilde{x}, \tilde{x}, \tilde{x}, \tilde{x})$. That is, the constant sequence $\{x_n\}$ with $x_n = \tilde{x}$ for all $n \geq -k \geq -l \geq \sigma$ is a solution of that equation.

Definition 2. Let $\tilde{x} \in (0, \infty)$ be an equilibrium point of Eq.(2). Then we have

* Corresponding author e-mail: mabdeltmoneam2014@yahoo.com

(i) An equilibrium point \tilde{x} of Eq.(2) is called locally stable if for every $\varepsilon > 0$ there exists $\delta > 0$ such that, if $x_{-\sigma}, \dots, x_{-l}, \dots, x_{-k}, \dots, x_{-1}, x_0 \in (0, \infty)$ with $|x_{-\sigma} - \tilde{x}| + \dots + |x_{-l} - \tilde{x}| + \dots + |x_{-k} - \tilde{x}| + \dots + |x_{-1} - \tilde{x}| + |x_0 - \tilde{x}| < \delta$, then $|x_n - \tilde{x}| < \varepsilon$ for all $n \geq -k \geq -l$.

(ii) An equilibrium point \tilde{x} of Eq.(2) is called locally asymptotically stable if it is locally stable and there exists $\gamma > 0$ such that, if $x_{-\sigma}, \dots, x_{-l}, \dots, x_{-k}, \dots, x_{-1}, x_0 \in (0, \infty)$ with $|x_{-\sigma} - \tilde{x}| + \dots + |x_{-l} - \tilde{x}| + \dots + |x_{-k} - \tilde{x}| + \dots + |x_{-1} - \tilde{x}| + |x_0 - \tilde{x}| < \gamma$, then

$$\lim_{n \rightarrow \infty} x_n = \tilde{x}.$$

(iii) An equilibrium point \tilde{x} of Eq.(2) is called a global attractor if for every $x_{-\sigma}, \dots, x_{-l}, \dots, x_{-k}, \dots, x_{-1}, x_0 \in (0, \infty)$ we have

$$\lim_{n \rightarrow \infty} x_n = \tilde{x}.$$

(iv) An equilibrium point \tilde{x} of Eq.(2) is called globally asymptotically stable if it is locally stable and a global attractor.

(v) An equilibrium point \tilde{x} of Eq.(2) is called unstable if it is not locally stable.

Definition 3. A sequence $\{x_n\}_{n=-\sigma}^{\infty}$ is said to be periodic with period r if $x_{n+r} = x_n$ for all $n \geq -\sigma$. A sequence $\{x_n\}_{n=-\sigma}^{\infty}$ is said to be periodic with prime period r if r is the smallest positive integer having this property.

Definition 4. Eq.(2) is called permanent and bounded if there exists numbers m and M with $0 < m < M < \infty$ such that for any initial conditions $x_{-\sigma}, \dots, x_{-l}, \dots, x_{-k}, \dots, x_{-1}, x_0 \in (0, \infty)$ there exists a positive integer N which depends on these initial conditions such that

$$m \leq x_n \leq M \quad \text{for all } n \geq N.$$

Definition 5. The linearized equation of Eq.(2) about the equilibrium point \tilde{x} is defined by the equation

$$z_{n+1} = \rho_0 z_n + \rho_1 z_{n-k} + \rho_2 z_{n-l} + \rho_3 z_{n-\sigma} = 0, \quad (3)$$

where

$$\rho_0 = \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x}, \tilde{x})}{\partial x_n}, \quad \rho_1 = \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x}, \tilde{x})}{\partial x_{n-k}},$$

$$\rho_2 = \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x}, \tilde{x})}{\partial x_{n-l}}, \quad \rho_3 = \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x}, \tilde{x})}{\partial x_{n-\sigma}}.$$

The characteristic equation associated with Eq.(3) is

$$\rho(\lambda) = \lambda^{\sigma+1} - \rho_0 \lambda^{\sigma} - \rho_1 \lambda^{\sigma-k} - \rho_2 \lambda^{\sigma-l} - \rho_3 = 0. \quad (4)$$

Theorem 1.[11]. Assume that F is a C^1 -function and let \tilde{x} be an equilibrium point of Eq.(2). Then the following statements are true.

(i) If all roots of Eq.(4) lie in the open unit disk $|\lambda| < 1$, then the equilibrium point \tilde{x} is locally asymptotically stable.

(ii) If at least one root of Eq.(4) has absolute value greater than one, then the equilibrium point \tilde{x} is unstable.

(iii) If all roots of Eq.(4) have absolute value greater than one, then the equilibrium point \tilde{x} is a source.

Theorem 2.[17]. Assume that ρ_0, ρ_1, ρ_2 and $\rho_3 \in \mathbb{R}$. Then

$$|\rho_0| + |\rho_1| + |\rho_2| + |\rho_3| < 1, \quad (5)$$

is a sufficient condition for the asymptotic stability of Eq.(2).

Theorem 3.[11]. Consider the difference equation (2). Let $\tilde{x} \in I$ be an equilibrium point of Eq.(2). Suppose also that

(i) F is a nondecreasing function in each of its arguments.

(ii) The function F satisfies the negative feedback property

$$[F(x, x, x, x) - x](x - \tilde{x}) < 0 \quad \text{for all } x \in I - \{\tilde{x}\},$$

where I is an open interval of real numbers. Then \tilde{x} is global attractor for all solutions of Eq.(2).

2 The local stability of the solutions

The equilibrium point \tilde{x} of Eq.(1) is the positive solution of the equation

$$\tilde{x} = (A + B + C + D)\tilde{x} + \frac{(b+h)\tilde{x}}{(d+e)\tilde{x}}. \quad (6)$$

If $0 < A + B + C + D < 1$, then the only positive equilibrium point \tilde{x} of Eq.(1) is given by

$$\tilde{x} = \frac{b+h}{(d+e)[1-(A+B+C+D)]}. \quad (7)$$

Let us now introduce a continuous function $F : (0, \infty)^4 \rightarrow (0, \infty)$ which is defined by

$$F(u_0, u_1, u_2, u_3) = Au_0 + Bu_1 + Cu_2 + Du_3 + \frac{bu_1 + hu_2}{du_1 + eu_2}. \quad (8)$$

Consequently, we get

$$\begin{cases} \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x})}{\partial u_0} = A = \rho_0, \\ \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x})}{\partial u_1} = B + \frac{(be-dh)[1-(A+B+C+D)]}{(d+e)(b+h)} = \rho_1, \\ \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x})}{\partial u_2} = C - \frac{(be-dh)[1-(A+B+C+D)]}{(d+e)(b+h)} = \rho_2, \\ \frac{\partial F(\tilde{x}, \tilde{x}, \tilde{x})}{\partial u_3} = D = \rho_3. \end{cases} \quad (9)$$

Thus, the linearized equation of Eq.(1) about \tilde{x} takes the form

$$z_{n+1} - \rho_0 z_n - \rho_1 z_{n-k} - \rho_2 z_{n-l} - \rho_3 z_{n-\sigma} = 0, \quad (10)$$

where ρ_0, ρ_1, ρ_2 and ρ_3 are given by (9).

Theorem 4.(i) If $be - dh = 0$, then the equilibrium point \tilde{x} of Eq.(2) is locally asymptotically stable.

(ii) If $be - dh \neq 0$, then equilibrium point \tilde{x} of Eq.(2) is locally asymptotically stable if the following condition hold:

$$\begin{aligned} &|B(d+e)(b+h) + (be-dh)[1-(A+B+C+D)]| \\ &+ |C(d+e)(b+h) - (be-dh)[1-(A+B+C+D)]| \\ &< (d+e)(b+h)[1-(A+B)]. \end{aligned} \quad (11)$$

Proof.(i) If $be - dh = 0$, we obtain

$$|\rho_0| + |\rho_1| + |\rho_2| + |\rho_3| = A + D + B + C < 1,$$

then we deduce from Theorem 2 that \tilde{x} of Eq.(2) is locally asymptotically stable.

(ii) If $be - dh \neq 0$, we deduce from (9) and (11) that $|\rho_0| + |\rho_1| + |\rho_2| + |\rho_3| < 1$, and hence \tilde{x} of Eq.(2) is locally asymptotically stable. Thus, the proof is now completed. \square

3 Periodic solutions

In this section, we study the existence of periodic solutions of Eq.(1). The following theorem states the necessary and sufficient conditions that the equation (1) has periodic solutions of prime period two.

Theorem 5. If k, l and σ are all even positive integers, then Eq.(1) has no prime period two solution.

Proof. Assume that there exists distinct positive solutions

$$\dots, P, Q, P, Q, \dots$$

of prime period two of Eq.(1). If k, l and σ are all even positive integers, then $x_n = x_{n-k} = x_{n-l} = x_{n-\sigma}$. It follows from Eq.(1) that

$$P = (A + B + C + D)Q + \frac{(b+h)}{(d+e)}, \quad (12)$$

and

$$Q = (A + B + C + D)P + \frac{(b+h)}{(d+e)}. \quad (13)$$

By subtracting (13) from (12), we get

$$(P - Q)[A + B + C + D + 1] = 0.$$

Since $A + B + C + D + 1 \neq 0$, then $P = Q$. This is a contradiction. Thus, the proof is now completed. \square

Theorem 6. If k, l and σ are all odd positive integers and $0 < A + B + C + D < 1$, then Eq.(1) has no prime period two solution.

Proof. If k, l and σ are all odd positive integers, then $x_{n+1} = x_{n-k} = x_{n-l} = x_{n-\sigma}$. It follows from Eq.(1) that

$$P = AQ + (B + C + D)P + \frac{(b+h)}{(d+e)}, \quad (14)$$

and

$$Q = AP + (B + C + D)Q + \frac{(b+h)}{(d+e)}. \quad (15)$$

By subtracting (15) from (14), we get

$$(P - Q)[(A + 1) - (B + C + D)] = 0.$$

Since $A + B + C + D < 1$, then $(A + 1) - (B + C + D) > 2A > 0$, hence $P = Q$. This is a contradiction. Thus, the proof is now completed. \square

Theorem 7. If k, l are even and σ is odd positive integers and $0 < A + B + C + D < 1$, then Eq.(1) has no prime period two solution.

Proof. If k, l are even and σ is odd positive integers, then $x_n = x_{n-k} = x_{n-l}$ and $x_{n+1} = x_{n-\sigma}$. It follows from Eq.(1) that

$$P = (A + B + C)Q + DP + \frac{(b+h)}{(d+e)}, \quad (16)$$

and

$$Q = (A + B + C)P + DQ + \frac{(b+h)}{(d+e)}. \quad (17)$$

By subtracting (17) from (16), we get

$$(P - Q)[(A + B + C + 1) - D] = 0,$$

Since $A + B + C + D < 1$, then $(A + B + C + 1) - D \neq 0$, hence $P = Q$. This is a contradiction. Thus, the proof is now completed. \square

Theorem 8. If σ is even and k, l are odd positive integers and $0 < A + B + C + D < 1$, then Eq.(1) has no prime period two solution.

Proof. If σ is even and k, l are odd positive integers, then $x_n = x_{n-\sigma}$ and $x_{n+1} = x_{n-k} = x_{n-l}$. It follows from Eq.(1) that

$$P = (A + D)Q + (B + C)P + \frac{(b + h)}{(d + e)}, \quad (18)$$

and

$$Q = (A + D)P + (B + C)Q + \frac{(b + h)}{(d + e)}. \quad (19)$$

By subtracting (19) from (18), we get

$$(P - Q)[(A + D + 1) - (B + C)] = 0,$$

Since $A + B + C + D < 1$, then $(A + D + 1) - (B + C) \neq 0$, hence $P = Q$. This is a contradiction. Thus, the proof is now completed. \square

Theorem 9. If k is even and l, σ are odd positive integers, then Eq.(1) has prime period two solution if the condition

$$(h - b)(d - e)\{(A + B) + [1 - (C + D)]\} > 4\{hd(A + B) + be[1 - (C + D)]\}, \quad (20)$$

is valid, provided $d > e, h > b, (C + D) < 1$.

Proof. If k is even and l, σ are odd positive integers, then $x_n = x_{n-k}$ and $x_{n+1} = x_{n-l} = x_{n-\sigma}$. It follows from Eq.(1) that

$$P = (A + B)Q + (C + D)P + \frac{bQ + hP}{dQ + eP}, \quad (21)$$

and

$$Q = (A + B)P + (C + D)Q + \frac{bP + hQ}{dP + eQ}. \quad (22)$$

Consequently, we get

$$eP^2 + dPQ = d(A + B)Q^2 + e(A + B)PQ + (C + D)dPQ + e(C + D)P^2 + bQ + hP, \quad (23)$$

and

$$eQ^2 + dPQ = d(A + B)P^2 + e(A + B)PQ + (C + D)dPQ + e(C + D)Q^2 + bP + hQ. \quad (24)$$

By subtracting (24) from (23), we get

$$P + Q = \frac{h - b}{d(A + B) + e[1 - (C + D)]}, \quad (25)$$

provided $h > b, (C + D) < 1$. By adding (23) and (24), we obtain

$$PQ = \frac{(h - b)\{hd(A + B) + be[1 - (C + D)]\}}{(d - e)[M_1 + (A + B)]\{d(A + B) + eM_1\}^2}, \quad (26)$$

where $M_1 = (1 - (C + D))$, provided $d > e, h > b, (C + D) < 1$. Assume that P and Q are two positive distinct real roots of the quadratic equation

$$t^2 - (P + Q)t + PQ = 0. \quad (27)$$

Thus, we deduce that

$$(P + Q)^2 > 4PQ. \quad (28)$$

Substituting (25) and (26) into (28), we get the condition (20). Thus, the proof is now completed. \square

Theorem 10. If l is even and k, σ are odd positive integers, then Eq.(1) has prime period two solution if the condition

$$(b - h)(e - d)\{(A + C) + [1 - (B + D)]\} > 4\{eb(A + C) + dh[1 - (B + D)]\}, \quad (29)$$

is valid, provided $e > d, b > h, (B + D) < 1$.

Proof. If l is even and k, σ are odd positive integers, then $x_n = x_{n-l}$ and $x_{n+1} = x_{n-k} = x_{n-\sigma}$. It follows from Eq.(1) that

$$P = (A + C)Q + (B + D)P + \frac{bP + hQ}{dP + eQ}, \quad (30)$$

and

$$Q = (A + C)P + (B + D)Q + \frac{bQ + hP}{dQ + eP}. \quad (31)$$

Consequently, we get

$$P + Q = \frac{b - h}{d[1 - (B + D)] + e(A + C)}, \quad (32)$$

where $(B + D) < 1, b > h$.

$$PQ = \frac{(b - h)\{eb(A + C) + dh[1 - (B + D)]\}}{(e - d)\{(A + C) + M_2\}\{e(A + C) + dM_2\}^2}, \quad (33)$$

where $M_2 = (1 - (B + D))$, $(B + D) < 1$. Substituting (32) and (33) into (28), we get the condition (29). Thus, the proof is now completed. \square

Theorem 11. If l, σ are even and k is odd positive integers, then Eq.(1) has prime period two solution if the condition

$$(b - h)(e - d)\{(A + C + D) + (1 - B)\} > 4\{eb(A + C + D) + dh(1 - B)\}, \quad (34)$$

is valid, provided $e > d, b > h, B < 1$.

Proof. If l, σ are even and k is odd positive integers, then $x_n = x_{n-l} = x_{n-\sigma}$ and $x_{n+1} = x_{n-k}$. It follows from Eq.(1) that

$$P = (A + C + D)Q + BP + \frac{bP + hQ}{dP + eQ}, \quad (35)$$

and

$$Q = (A + C + D)P + BQ + \frac{bQ + hP}{dQ + eP}. \quad (36)$$

Consequently, we get

$$P + Q = \frac{b - h}{d(1 - B) + e(A + C + D)}, \quad (37)$$

where $b > h, B < 1$,

$$PQ = \frac{(b - h)\{e b (A + C + D) + dh (1 - B)\}}{(e - d)\{M_3 + (1 - B)\}\{e M_3 + d (1 - B)\}^2}, \quad (38)$$

where $M_3 = (A + C + D), e > d, b > h, B < 1$. Substituting (37) and (38) into (28), we get the condition (34). Thus, the proof is now completed. \square

Theorem 12. *If k, σ are even and l is odd positive integers, then Eq.(1) has prime period two solution if the condition*

$$(h - b)(d - e)\{(A + B + D) + (1 - C)\} > 4\{eb(1 - C) + dh(A + B + D)\}, \quad (39)$$

is valid, provided $d > e, h > b, C < 1$.

Proof. If k, σ are even and l is odd positive integers, then $x_n = x_{n-k} = x_{n-\sigma}$ and $x_{n+1} = x_{n-l}$. It follows from Eq.(1) that

$$P = (A + B + D)Q + CP + \frac{bQ + hP}{dQ + eP}, \quad (40)$$

and

$$Q = (A + B + D)P + CQ + \frac{bP + hQ}{dP + eQ}. \quad (41)$$

Consequently, we get

$$P + Q = \frac{h - b}{e(1 - C) + d(A + B + D)}, \quad (42)$$

where $h > b, C < 1$,

$$PQ = \frac{(h - b)\{dh(A + B + D) + e b (1 - C)\}}{(d - e)\{(1 - C) + M_4\}\{d M_4 + e (1 - C)\}^2}, \quad (43)$$

where $M_4 = (A + B + D), d > e, h > b, C < 1$. Substituting (42) and (43) into (28), we get the condition (39). Thus, the proof is now completed. \square

4 Boundedness of the solutions

In this section, we investigate the boundedness of the positive solutions of Eq.(1).

Theorem 13. *If $0 < A + B + C + D < 1$, then every solution of Eq.(1) is bounded from above by the quantity*

$$x_0 + \left(1 + \sum_{i=1}^{n-1} A^i\right) \left(\frac{b e + dh}{de}\right),$$

where $A < 1$, and x_0 is an initial condition.

Proof. Let $\{x_n\}_{n=-\sigma}^{\infty}$ be a solution of Eq.(1). It follows from Eq.(1) that

$$\begin{aligned} x_{n+1} &= Ax_n + Bx_{n-k} + Cx_{n-l} + Dx_{n-\sigma} + \frac{bx_{n-k} + hx_{n-l}}{dx_{n-k} + ex_{n-l}} \\ &\leq Ax_n + Bx_{n-k} + Cx_{n-l} + Dx_{n-\sigma} + \frac{b}{d} + \frac{h}{e}. \end{aligned}$$

Since the initial conditions satisfy the inequality $x_{-\sigma} < \dots < x_{-l} < \dots < x_{-k} < \dots < x_{-1} < x_0$, then we get

$$x_1 \leq (A + B + C + D)x_0 + \left(\frac{b}{d} + \frac{h}{e}\right),$$

and hence

$$x_1 \leq x_0 + \left(\frac{be + hd}{de}\right). \quad (44)$$

With the aid of (44) we have

$$\begin{aligned} x_2 &\leq Ax_1 + Bx_{1-k} + Cx_{1-l} + Dx_{1-\sigma} + \left(\frac{be + hd}{de}\right) \\ &\leq (A + B + C + D)x_0 + (A + 1) \left(\frac{be + hd}{de}\right), \end{aligned}$$

and hence

$$x_2 \leq x_0 + (A + 1) \left(\frac{be + hd}{de}\right). \quad (45)$$

With the aid of (45) we have

$$\begin{aligned} x_3 &\leq Ax_2 + Bx_{2-k} + Cx_{2-l} + Dx_{2-\sigma} + \left(\frac{be + hd}{de}\right) \\ &\leq (A + B + C + D)x_0 + (A + 1) \left(\frac{be + hd}{de}\right), \end{aligned}$$

and hence

$$x_3 \leq x_0 + [A^2 + A + 1] \left(\frac{be + hd}{de}\right). \quad (46)$$

By continuing this process, we deduce that

$$x_n \leq x_0 + \left(1 + \sum_{i=1}^{n-1} A^i\right) \left(\frac{b e + dh}{de}\right), \quad n = 1, 2, 3, \dots \quad (47)$$

where we define $\sum_{i=1}^0 A^i = 0$. Then every solution of Eq.(1) is bounded from above by the quantity

$$x_0 + \left(1 + \sum_{i=1}^{n-1} A^i\right) \left(\frac{b e + dh}{de}\right).$$

Thus, the proof is now completed. \square

5 Global stability

In this section, we study the global asymptotic stability of the positive solutions of Eq.(1).

Theorem 14. If $0 < A + B + C + D < 1$, then the equilibrium point \tilde{x} given by (7) of Eq.(1) is global attractor.

Proof. We consider the following function

$$F(x, y, z, w) = Ax + By + Cz + Dw + \frac{by + hz}{dy + ez}. \quad (48)$$

It is easy to verify the condition (i) of Theorem 3. Let us now verify the condition (ii) of Theorem 3 as follows:

$$[F(x, x, x, x) - x](x - \tilde{x}) = \left\{ (A + B + C + D)x + \frac{b+h}{d+e} - x \right\} \\ \times \left\{ x - \frac{b+h}{(d+e)[1 - (A+B+C+D)]} \right\}$$

$$= - \left\{ [1 - (A + B + C + D)]x - \frac{b+h}{d+e} \right\} \\ \times \left\{ x - \frac{b+h}{(d+e)[1 - (A+B+C+D)]} \right\} \\ = - [1 - (A + B + C + D)] \\ \times \left\{ x - \frac{b+h}{(d+e)[1 - (A+B+C+D)]} \right\}^2. \quad (49)$$

Since $0 < A + B + C + D < 1$, then we deduce from (49) that

$$[F(x, x, x, x) - x](x - \tilde{x}) < 0. \quad (50)$$

According to Theorem 3, \tilde{x} is global attractor. Thus, the proof is now completed. \square


On combining the two Theorems 4 and 14, we have the following result:

Theorem 15. The equilibrium point \tilde{x} given by (7) of Eq.(1) is globally asymptotically stable.


6 Numerical examples

In order to illustrate the results of the previous section and to support our theoretical discussions, we consider some numerical examples in this section. These examples represent different types of qualitative behavior of solutions of Eq.(1).


Example 1. Figure 1, shows that Eq.(1) has no prime period two solutions if $k = 2$, $l = 4$, $\sigma = 6$, $x_{-6} = 1$, $x_{-5} = 2$, $x_{-4} = 3$, $x_{-3} = 4$, $x_{-2} = 5$, $x_{-1} = 6$, $x_0 = 7$, $A = 300$, $B = 200$, $C = 100$, $D = 75$, $b = 50$, $h = 40$, $d = 30$, $e = 20$.




Example 2. Figure 2, shows that Eq.(1) has no prime period two solutions if $k = 1$, $l = 3$, $\sigma = 5$, $x_{-5} = 1$, $x_{-4} = 2$, $x_{-3} = 3$, $x_{-2} = 4$, $x_{-1} = 5$, $x_0 = 6$, $A = 75$, $B = 50$, $C = 25$, $D = 20$, $b = 5$, $h = 4$, $d = 3$, $e = 2$.




Example 3. Figure 3, shows that Eq.(1) has no prime period two solutions if $k = 2$, $l = 4$, $\sigma = 5$, $x_{-5} = 1$, $x_{-4} = 2$, $x_{-3} = 3$, $x_{-2} = 4$, $x_{-1} = 5$, $x_0 = 6$, $A = 10$, $B = 20$, $C = 30$, $D = 40$, $b = 5$, $h = 4$, $d = 3$, $e = 2$.



Example 4. Figure 4, shows that Eq.(1) has no prime period two solutions if $k = 1, l = 3, \sigma = 4, x_{-4} = 1, x_{-3} = 2, x_{-2} = 3, x_{-1} = 4, x_0 = 5, A = 100, B = 50, C = 25, D = 200, b = 5, h = 4, d = 3, e = 2.$



Example 5. shows that Eq.(1) is globally asymptotically stable if $k = 1, l = 3, \sigma = 4, x_{-4} = 1, x_{-3} = 2, x_{-2} = 3, x_{-1} = 4, x_0 = 5, A = 0.01, B = 0.02, C = 0.03, D = 0.04, b = 5, h = 4, d = 3, e = 2000.$



7 Conclusion

We have discussed some properties of the nonlinear rational difference equation (1), namely the periodicity, the boundedness and the global stability of the positive solutions of this equation. We gave some figures to illustrate the behavior of these solutions. Our results in this article can be considered as a more generalization than the results obtained in Refs.[1,26,30,31]. Note that example 1 verifies Theorem 5 which shows that if k, l and σ are all even positive integers, then Eq.(1) has no prime period two solution and example 2 verifies Theorem 6

which shows that if k, l and σ are all odd positive integers, then Eq.(1) has no prime period two solution and example 3 verifies Theorem 7 which shows that if k, l are even and σ is odd positive integers, then Eq.(1) has no prime period two solution. But example 4 verifies Theorem 8 which shows that if σ is even and k, l are odd positive integers, then Eq.(1) has no prime period two solution, while example 5 verifies Theorem 15 which shows that Eq.(1) is globally asymptotically stable.

Acknowledgement

The authors wish to thank the referees for their suggestions and comments.

References

- [1] E. M. Elabbasy, H. El- Metwally and E. M. Elsayed, On the difference equation $x_{n+1} = ax_n - bx_n / (cx_n - dx_{n-1})$, *Advances in Difference Equations*, **2006**, 1-10.
- [2] H. El-Metwally and E. M. Elsayed, Solution and behavior of a third rational difference equation, *Utilitas Mathematica*, **88**, 27-42 (2012).
- [3] M. A. El-Moneam and S.O. Alamoudy, On study of the asymptotic behavior of some rational difference equations, *DCDIS Series A: Mathematical Analysis* (to appear).
- [4] E. M. Elsayed, Solution and attractivity for a rational recursive sequence, *Discrete Dynamics in Nature and Society*, **2011**, 17 pages.
- [5] E. M. Elsayed, Solutions of rational difference system of order two, *Mathematical and Computer Modelling*, **55**, 378-384 (2012).
- [6] E. M. Elsayed, Behavior and expression of the solutions of some rational difference equations, *J. Computational Analysis and Applications*, **15**, 73-81 (2013).
- [7] E. M. Elsayed and M. M. El-Dessoky, Dynamics and behavior of a higher order rational recursive sequence, *Advances in Difference Equations*, **2012**, (2012).
- [8] E. M. Elsayed and H. A. El-Metwally, On the solutions of some nonlinear systems of difference equations, *Advances in Difference Equations*, **2013**, (2013).
- [9] M. E. Erdogan, C. Cinar and I. Yalcinkaya, On the dynamics of the recursive sequence, *Computers & Mathematics with Applications*, **61**, 533-537 (2011).
- [10] M. E. Erdogan, C. Cinar and I. Yalcinkaya, On the dynamics of the recursive sequence, *Mathematical and Computer Modelling*, **54**, 1481-1485 (2011).
- [11] E. A. Grove and G. Ladas, Periodicities in nonlinear difference equations, *Chapman & Hall / CRC*, **4**, (2005).
- [12] T. F. Ibrahim, Solutions of a system of nonlinear fractional recursion equations, *International Journal of Mathematical Education: Policy and Practice*, **2**, 49-54 (2012).
- [13] T. F. Ibrahim, Periodicity and solution of rational recurrence relation of order six, *Applied Mathematics*, **3**, 729-733 (2012).
- [14] T. F. Ibrahim, Three-dimensional max-type cyclic system of difference equations, *International Journal of Physical Sciences*, **8**, 629-634 (2013).

- [15] T. F. Ibrahim and N. Touafek, On a third-order rational difference equation with variable coefficients, *DCDIS Series B: Applications & Algorithms*, **20**, 251-264 (2013).
- [16] V. L. Kocic and G. Ladas, *Global behavior of nonlinear difference equations of higher order with applications*, Kluwer Academic Publishers, Dordrecht, (1993).
- [17] M. R. S. Kulenovic and G. Ladas, *Dynamics of second order rational difference equations with open problems and conjectures*, Chapman & Hall / CRC, Florida, (2001).
- [18] N. Touafek and E. M. Elsayed, On the solutions of systems of rational difference equations, *Mathematical and Computer Modelling*, **55**, 1987-1997 (2012).
- [19] N. Touafek and E. M. Elsayed, On the periodicity of some systems of nonlinear difference equations, *Bull. Math. Soc. Sci. Math. Roumanie, Tome*, **55**, 217-224 (2012).
- [20] I. Yalcinkaya and C. Cinar, On the dynamics of the difference equation $x_{n+1} = ax_{n-k}/(b + cx_n^p)$, *Fasc. Math.*, **42**, 141-148 (2009).
- [21] E. M. E. Zayed, On the rational recursive sequence $x_{n+1} = Ax_n + Bx_{n-k} + (1 + x_n x_{n-k}) / (px_n + x_{n-k})$, *Acta Math. Vietnamica*, **37**, 251-266 (2012).
- [22] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = (D + \alpha x_n + \beta x_{n-1} + \gamma x_{n-2}) / (Ax_n + Bx_{n-1} + Cx_{n-2})$, *Comm. Appl. Nonl. Anal.*, **12**, 15-28 (2005).
- [23] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = (\alpha x_n + \beta x_{n-1} + \gamma x_{n-2} + \delta x_{n-3}) / (Ax_n + Bx_{n-1} + Cx_{n-2} + Dx_{n-3})$, *J. Appl. Math. & Computing*, **22**, 247-262 (2006).
- [24] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = (A + \sum_{i=0}^k \alpha_i x_{n-i}) / \sum_{i=0}^k \beta_i x_{n-i}$, *Math. Bohemica*, **133**, 225-239 (2008).
- [25] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = (A + \sum_{i=0}^k \alpha_i x_{n-i}) / (B + \sum_{i=0}^k \beta_i x_{n-i})$, *Int. J. Math. & Math. Sci.*, **2007**, 12 pages.
- [26] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = ax_n - bx_n / (cx_n - dx_{n-k})$, *Comm. Appl. Nonlinear Analysis*, **15**, 47-57 (2008).
- [27] E. M. E. Zayed and M. A. El-Moneam, On the Rational Recursive Sequence $x_{n+1} = (\alpha + \beta x_{n-k}) / (\gamma - x_n)$, *J. Appl. Math. & Computing*, **31**, 229-237 (2009).
- [28] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = Ax_n + (\beta x_n + \gamma x_{n-k}) / (Cx_n + Dx_{n-k})$, *Comm. Appl. Nonl. Anal.*, **16**, 91-106 (2009).
- [29] E. M. E. Zayed and M. A. El-Moneam, On the Rational Recursive Sequence $x_{n+1} = \gamma x_{n-k} + (ax_n + bx_{n-k}) / (cx_n - dx_{n-k})$, *Bull. Iranian Math. Soc.*, **36**, 103-115 (2010).
- [30] E. M. E. Zayed and M. A. El-Moneam, On the global attractivity of two nonlinear difference equations, *J. Math. Sci.*, **177**, 487-499 (2011).
- [31] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive two sequences $x_{n+1} = ax_{n-k} + bx_{n-k} / (cx_n + \delta dx_{n-k})$, *Acta Math. Vietnamica*, **35**, 355-369 (2010).
- [32] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = Ax_n + Bx_{n-k} + (\beta x_n + \gamma x_{n-k}) / (Cx_n + Dx_{n-k})$, *Acta Appl. Math.*, **111**, 287-301 (2010).
- [33] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = (A + \alpha_0 x_n + \alpha_1 x_{n-\sigma}) / (B + \beta_0 x_n + \beta_1 x_{n-\tau})$, *Acta Math. Vietnamica*, **36**, 73-87 (2011).
- [34] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = (\alpha_0 x_n + \alpha_1 x_{n-l} + \alpha_2 x_{n-k}) / (\beta_0 x_n + \beta_1 x_{n-l} + \beta_2 x_{n-k})$, *Math. Bohemica*, **135**, 319-336 (2010).
- [35] E. M. E. Zayed and M. A. El-Moneam, On the global asymptotic stability for a rational recursive sequence, *Iranian Journal of science and Technology (A: sciences)*, **35**, 333-339 (2011).
- [36] E. M. E. Zayed and M. A. El-Moneam, On the rational recursive sequence $x_{n+1} = \frac{\alpha_0 x_n + \alpha_1 x_{n-l} + \alpha_2 x_{n-m} + \alpha_3 x_{n-k}}{\beta_0 x_n + \beta_1 x_{n-l} + \beta_2 x_{n-m} + \beta_3 x_{n-k}}$, *WSEAS Transactions Math.*, **11**, 373-382 (2012).
- [37] E. M. E. Zayed and M. A. El-Moneam, On the qualitative study of the nonlinear difference equation $x_{n+1} = \frac{\alpha x_{n-\sigma}}{\beta + \gamma x_{n-\tau}}$, *Fasciculi Mathematici*, **50**, 137-147 (2013).



M. A. El-Moneam
 Assistant Professor of Mathematics (from 2010 till now) at Mathematics Department, Faculty of Science and Arts in Farasan, Jazan University, Kingdom of Saudi Arabia. His interests are the nonlinear differential equations, Nonlinear

difference equations. He Published many articles in famous International Journals around the world. He has reviewed many articles for many international Journals. He has got MSC degree in Mathematics from Faculty of Science, at Zagazig University, Egypt, supervised Prof. E. M. E. Zayed. He has got his PHD in Mathematics from Faculty of Science, at Zagazig University, Egypt, supervised Prof. E. M. E. Zayed. He has got the Man of the Year award, (2011), by the American Biographical, Institute, U.S.A. He is a member of International Society of Difference Equations (ISDE) in U.S.A. He is a member of peer reviewer in AIP Conference Proceeding 2010 Athena (2010 , 2011, 2012, 2013). He is a member of peer reviewer in the 2nd Scientific Conference for students of higher education in Jeddah, Kingdom of Saudi Arabia (1432 G.). He is a member of peer reviewer in the 3rd Scientific Conference for students of higher education in Al Khobar, Kingdom of Saudi Arabia 1433. He is a member of peer reviewer in the 4th Scientific Conference for students of higher education in Makkah, Kingdom of Saudi Arabia 1434. He is a peer reviewed to the Scientific research No. (164/662/1432), King Abd Al Aziz University, Kingdom of Saudi Arabia (1432 H.). He edited and reviewed the Scientific book No. (208), Taif University, Kingdom of Saudi Arabia, 2013 G., (1434 H.). He has attended the Scientific sessions of the 4th international Exhibition and Conference on Higher Education (16-17/4/2013 G.), Riyadh, Kingdom of Saudi Arabia. He has attended the 1st Saudi Scientific Publishing Conference (27-29/3/1435 H.)(28-30/1/2014 G.) King Khalid University, Abha, Kingdom of Saudi Arabia.



E. M. E. Zayed
 Professor of Mathematics (from 1989 till now) at Mathematics Department, Faculty of Science, Zagazig University, Egypt. His interests are the nonlinear PDEs in Mathematical Physics, Inverse problems in differential equations,

Nonlinear difference equations. He Published about 225 articles in famous International Journals around the world. He was the head of Mathematics Department at Zagazig University, Egypt from 2001 till 2006. He is an Editor of WSEAS Transaction on Mathematics. He has got some Mathematical prizes by the Egyptian Academy of scientific research and Technology. He got the Medal of Science and Arts of the first class from the president of Egypt. He got the Medal of Excellent of the first class from the president of Egypt. He is one of the Editorial Board of the Punjab University Journal of Mathematics. He have reviewed many articles for many international Journals . He has got his BSC of Mathematics from Tanta University, Egypt 1973 . He has got two MSC degrees in Mathematics .The first one from Al-Azher University, Egypt 1977, while the second one from Dundee University,Scotland, UK 1978. He has got his PHD in Mathematics from Dundee University, Scotland, UK 1981. He has got the Man of the Year award, (1993), by the American Biographical, Institute, U.S.A. He has got the Research Fellow Medal, (1993), by the American Biographical Institute, U.S.A. The Curriculum Vitae of Professor E.M.E.Zayed has been published by Men of achievement of the International Biographical center, Cambridge, vol. 16, (1995), p.534 The Curriculum Vitae of Professor E.M.E.Zayed has been published by Dictionary of International Biography of the International Biographical center, Cambridge, vol. 23, (1995), p.714. He was a Professor of Mathematics at Taif University, Saudi Arabia (2006- 2010).